


# Genital psoriasis: clinical, epidemiological and therapeutic aspects

## *Psoríase genital: aspectos clínicos, epidemiológicos e terapêuticos*

Daiane de Souza<sup>1</sup>\*, João P. Fabrini-da Silva<sup>1</sup>, Rogério Nabor-Kondo<sup>2</sup>, and Cássio R. Moreira<sup>1</sup>

<sup>1</sup>Autarquia Municipal de Saúde de Apucarana, Apucarana; <sup>2</sup>Universidade Estadual de Londrina, Londrina. Brazil

### Abstract

Genital psoriasis presents as erythematous plaques with variable scaling and symptoms such as pruritus, burning, and dyspareunia, which can significantly affect the quality of life, psychological well-being, and sexual health of affected patients. It is estimated that approximately 1.2-2.5% of the population has psoriasis; however, genital involvement is not widely described and may be underestimated, as these areas are often not actively questioned or assessed during the clinical examination. This article aims to explore the clinical aspects of genital psoriasis, including its prevalence, impact on sexual health, and treatment options.

**Keywords:** Genital diseases. Genital psoriasis. Psoriasis. Therapy.

### Resumo

A psoríase genital manifesta-se como placas eritematosas com descamação variável e sintomas como prurido, ardor e dispareunia, que podem afetar significativamente a qualidade de vida, o bem-estar psicológico e a saúde sexual dos pacientes acometidos. Estima-se que aproximadamente 1.2% a 2.5% da população apresente psoríase; contudo, o acometimento genital não é amplamente descrito e pode ser subestimado, uma vez que essas áreas frequentemente não são ativamente investigadas ou avaliadas durante o exame clínico. Este artigo tem como objetivo explorar os aspectos clínicos da psoríase genital, incluindo sua prevalência, impacto na saúde sexual e opções terapêuticas.

**Palavras-chave:** Doenças genitais. Psoríase genital. Psoríase. Terapêutica.

#### \*Correspondence:

Daiane de Souza

E-mail: daiane\_souzab@hotmail.com

2795-501X / © 2026 Portuguese Society of Dermatology and Venereology. Published by Permayer. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Received: 20-12-2025

Accepted: 30-01-2026

DOI: 10.24875/PJDV.25000100

Available online: 12-03-2026

Port J Dermatol and Venereol. 2026;84(2):79-85

[www.portuguesejournalofdermatology.com](http://www.portuguesejournalofdermatology.com)

## Introduction

Psoriasis is a chronic, immune-mediated inflammatory disease characterized by erythematous, scaly, infiltrated, and pruritic plaques on the skin. It typically affects the extensor surfaces, scalp, palms, soles, and nails; however, it can involve any area of the integument. It has a negative impact on the quality of life of affected patients and can be associated with comorbidities such as psoriatic arthritis.<sup>1</sup>

Genital psoriasis usually presents as erythematous, well-defined plaques with varying degrees of desquamation. Symptoms such as pruritus, burning, and dyspareunia have been reported.<sup>2</sup> Therefore, genital psoriasis can significantly affect the quality of life, psychological well-being, and sexual health of affected patients.<sup>3</sup>

In Brazil, the estimated prevalence of psoriasis ranges from 1.2% to 2.5%.<sup>4,5</sup> However, genital involvement is not widely described. It is estimated that genital involvement occurs in more than two-thirds of patients;<sup>6</sup> however, this proportion may be underestimated, as these areas may not be actively questioned or assessed during the clinical examination.<sup>7</sup>

This paper aims to explore clinical aspects of genital psoriasis, including its prevalence, impact on sexual health and treatment options.

## Epidemiology

Among the various dermatoses that can affect the genitals, psoriasis is the most common.<sup>8</sup> A systematic review conducted by Meeuwis et al., including 18 studies and a total of 22,116 patients with psoriasis, showed that approximately 63% of patients will present some genital involvement during the course of the disease. This proportion increases to 79% when the patient has the flexural or inverted clinical form of the disease.<sup>6</sup>

In a prospective study totaling 776 patients with psoriasis, 43.2% of the patients had genital involvement.<sup>2</sup> An Indian study examining 852 patients with psoriasis reported a lower prevalence of 11.7%.<sup>9</sup> A prospective observational study with a total of 262 patients, 42% Chinese, 36.6% Malay and 21.4% Indian or other nationalities, showed 46.1% genital involvement by psoriasis currently or previously to the study.<sup>7</sup> Curiously, this study also identified that being Chinese was a risk factor for genital psoriasis when compared to the other ethnicities included.<sup>7</sup>

Men are more frequently affected by genital psoriasis than women<sup>6,7,10</sup> and it manifests simultaneously with

lesions on other parts of the body in the majority of cases, with isolated genital involvement being reported in only 2-5% of cases.<sup>11</sup> More severe cases, as assessed by the psoriasis area and severity index (PASI), and the greater use of systemic therapies have been associated with genital psoriasis.<sup>2</sup> A Swiss study evaluated 109 children with psoriasis, and genital involvement was present in around 40% of them and was often the first manifestation of the disease.<sup>12</sup> Circumcision apparently does not influence the development of genital psoriasis in men.<sup>2</sup>

## Clinical manifestations


The most characteristic lesions are well-defined erythematous plaques with minimal infiltration. Unlike lesions in other regions of the body, desquamation may be absent, possibly due to increased local humidity and friction.<sup>13</sup> However, typical scales may be present in more keratinized areas (Fig. 1).<sup>14</sup> The areas mostly affected in men include the body of the penis, the scrotum, and, less commonly, the glans (Fig. 2). In women, the labia majora are the most commonly affected areas and are often symmetrically involved, followed by the perineum.<sup>2</sup>

Pruritus is the main symptom of genital psoriasis, reported by more than 86% of patients; however, around half of the cases experience pain, burning, and discomfort during sexual intercourse due to the presence of fissures.<sup>2</sup> Due to the Koebner phenomenon, genital psoriasis can be aggravated by irritation caused by urine and feces, tight clothing, and trauma from sexual intercourse.<sup>15</sup>


Genital psoriasis is not a static disease; in general, its worsening occurs in parallel with the progression of non-genital lesions, although there are exceptions.<sup>16</sup> The main worsening factors traditionally described include emotional stress, trauma, some medications, and friction (related to sexual activity or wearing tight clothing).<sup>3</sup> Studies show that up to 34% of patients experience post-coital worsening.<sup>2,17</sup>

The diagnosis of genital psoriasis is essentially clinical, and biopsy is avoided in these cases due to the sensitivity of the area. Therefore, as a non-invasive method, dermoscopy may serve as an additional diagnostic tool. Dotted vessels and dilated, tortuous capillaries distributed over a homogeneous reddish background represent the most frequently described pattern.<sup>18</sup>

Psoriasis is a condition that generally does not lead to tissue fibrosis; however, Albert et al. described two cases of female patients with long-term genital psoriasis who developed fibrosis and loss of the labia minora.<sup>19</sup>



**Figure 1.** Erythematous and scaly plaques affecting the base of the penis and the pubic region of a male patient with psoriasis.



**Figure 2.** Erythematous plaques affecting the dorsal region of the penile body in a male patient with psoriasis.

Secondary infection by bacteria, fungi, or viruses is relatively common and can aggravate symptoms or cause atypical manifestations. Therefore, in the event of clinical suspicion, the infection should be treated promptly.<sup>13</sup> A retrospective observational study showed that more than a third of patients with genital psoriasis had positive swabs for *Candida albicans* or *Staphylococcus* spp.<sup>20</sup>

## Differential diagnoses

The differential diagnosis of genital psoriasis encompasses a broad spectrum of inflammatory and infectious dermatoses, including candidiasis, contact dermatitis, lichen planus, lichen sclerosus, and sexually transmitted infections, all of which may exhibit overlapping clinical features in the genital region.<sup>11,21</sup> Genital psoriasis may pose a significant diagnostic challenge, particularly when lesions are isolated and not accompanied by extragenital manifestations.<sup>11,22</sup> In such circumstances, clinical overlap with premalignant and malignant conditions must be considered. Squamous cell carcinoma, including erythroplasia of Queyrat, may closely mimic genital or inverse psoriasis, potentially leading to diagnostic misinterpretation and delayed treatment.<sup>23</sup> Although the diagnosis of genital psoriasis is typically clinical, current evidence supports a low threshold for performing a skin biopsy in cases of solitary, persistent, atypical, or treatment-refractory genital lesions, to exclude malignancy and ensure diagnostic accuracy.<sup>11,22,23</sup>

## Severity assessment tools

The assessment tools traditionally used in psoriasis may not be suitable for the genital form of the disease. The PASI and body surface area are tools that depend on the area of the affected body surface, so in genital psoriasis alone, they will not be able to demonstrate the severity of the disease adequately.<sup>24</sup> Although the dermatology life quality index includes, in item 9, the impact of the disease on sexual life, it is not specific to genital psoriasis.<sup>25</sup> Therefore, more specific tools have been developed in order to refine this assessment, including the genital psoriasis sexual frequency questionnaire and the genital psoriasis symptoms scale.<sup>26</sup>

## Treatment

The treatment of genital psoriasis is challenging due to the particularities of the skin in this area. Patients should be informed that the disease is neither transmissible nor the result of poor hygiene. In addition, teaching them to use non-irritating cleansers, topical medication in sufficient quantities, and moisturizing the area helps with the therapeutic response. Furthermore, the use of intimate lubricants during sexual intercourse reduces friction and, consequently, the risk of Koebner's phenomenon.<sup>27</sup>

Therapy with topical corticosteroids, alone or in combination with systemic drugs, has been shown to be the initial treatment of choice for genital psoriasis.<sup>28</sup> As the skin is thinner and subject to natural occlusion, percutaneous absorption is increased;<sup>29</sup> therefore, the use of low-potency topical corticosteroids, such as hydrocortisone, is recommended, and their effectiveness has been demonstrated in prospective and retrospective studies.<sup>6,20,30,31</sup> It is recommended not to extrapolate the use time of four continuous weeks to minimize the risk of well-known adverse events, including skin atrophy, telangiectasias, and stretch marks.<sup>28</sup>

Although low or medium potency corticosteroids are preferable for sensitive areas such as the genitals, high potency corticosteroids may be indicated for short periods in moderate to severe cases or to induce an initial clinical response.<sup>20</sup> Formulations with Vitamin D analogues, such as calcipotriol, can be used in conjunction with corticosteroids or even after their use.<sup>31</sup>

Topical calcineurin inhibitors (TCI), such as tacrolimus and pimecrolimus, are alternatives for long-term topical therapies. They are associated with milder and more manageable adverse effects, as they do not interfere with collagen synthesis, so the risk of skin atrophy is significantly lower compared to corticosteroids.<sup>32</sup> However, they can cause pruritus and burning at the application site.

Although the indication for the treatment of psoriasis is not included in the package leaflet for TCIs, some studies have shown benefits. A study by Bissonnette et al. showed a satisfactory response and tolerability in a group of 12 men with genital psoriasis who used tacrolimus 0.1% ointment for 8 weeks.<sup>33</sup> A study of 10 patients with facial and genital psoriasis showed that tacrolimus 0.1% ointment improved the symptoms of genital psoriasis after 12 weeks of use.<sup>34</sup>

Topical Vitamin D analogs, such as calcitriol and calcipotriol, are another class of drugs that can be used for a long time in genital and inverted psoriasis. In general, studies show that both calcitriol and calcipotriol are safe, but less effective than corticosteroids or TCI.<sup>35</sup> A randomized direct comparison study between calcitriol 3 µg/g ointment and calcipotriol 50 µg/g ointment, both applied twice a day, showed that calcitriol was not only more effective, but also better tolerated in the treatment of inverted psoriasis.<sup>36</sup>


In general, phototherapy is not indicated for the treatment of genital psoriasis due to the increased risk of skin cancer in this area. A prospective study of 892 men with psoriasis exposed for a long time to psoralens and ultraviolet A (PUVA) radiation had an increased risk

of tumors in the penis and scrotum, particularly squamous cell carcinoma. Furthermore, even after controlling for levels of exposure to PUVA, high levels of exposure to ultraviolet B radiation were also associated with an increased risk of genital cancer.<sup>37</sup>

Up to now, there have been no large clinical trials evaluating the efficacy and safety of traditional oral systemic therapies for genital psoriasis. One study demonstrated successful treatment with methotrexate, but a second described only a partial response to the drug.<sup>12,20</sup> In addition, dapsone was used with a satisfactory response in two cases of pustular psoriasis affecting the skin of the penis.<sup>38,39</sup> On the other hand, a randomized, placebo-controlled study showed that apremilast, an oral phosphodiesterase-4 inhibitor, improved the quality of life of patients with psoriasis involving limited and spatial areas, including the genitals.<sup>40</sup> More specifically, another recent study evaluated the response to apremilast in patients with genital psoriasis, which also showed benefits.<sup>41</sup>

At present, the biologic agent with the specific indication for the treatment of genital psoriasis is ixekizumab, which is a high-affinity monoclonal antibody against interleukin (IL)-17A. Several studies have shown significant improvement in lesion appearance, pruritus, sexual health, and quality of life in patients with resistant genital psoriasis treated with ixekizumab.<sup>42-45</sup> In addition, a Phase III, randomized, placebo-controlled clinical study demonstrated long-term efficacy and safety up to week 52 of treatment with ixekizumab.<sup>44</sup> Patients received subcutaneous injections of 80 mg of ixekizumab (n = 74) or placebo (n = 74) every 2 weeks after an initial dose of 160 mg at week 0 until week 12; after which the study entered an open phase in which all patients received 80 mg every 4 weeks until week 52. As a result, in the initial phase of the study, 73% of the patients who received the drug achieved total or almost total improvement. Similarly, 79% of the patients who were initially in the placebo group achieved similar results at the end of week 52.

In addition to clinical trials, a real-life study involving 1978 patients with moderate to severe psoriasis, of whom around 25% had genital involvement, showed that in this subgroup of patients the IL-17A inhibitor biologics (ixekizumab and secukinumab) were significantly more effective than the other biologics agents used by the patients in the study (guselkumab, risankizumab, ustekinumab and adalimumab).<sup>46</sup> When compared individually with ixekizumab, ustekinumab, and guselkumab were the least effective of the biologics agents used.



**Figure 3.** Therapeutic flowchart for genital psoriasis.

Similarly, in a real-life study of 255 patients with psoriasis treated with adalimumab, etanercept, or ustekinumab, genital psoriasis was associated with complete resolution of the condition (PASI100) when patients were treated with adalimumab.<sup>47</sup> In this context, Orsini et al. showed satisfactory results in a real-life scenario in which patients with psoriasis were treated with risankizumab. Of a total of 202 patients, 72 had genital involvement. At the end of weeks 16, 28, and 52 of risankizumab use, 85%, 93% and 100% showed total or almost total improvement, respectively.<sup>48</sup>

Finally, the Italian GULLIVER study, a 12-week interim analysis, evaluated the efficacy and safety of guselkumab, a fully human monoclonal antibody targeting the p19 subunit of IL-23, in the treatment of genital and facial psoriasis. With 351 patients included, 204 of whom had genital psoriasis, it was observed that 76.5% of patients achieved an static physician's global assessment score of 0 (clean) or 1 (almost clean) in the genital

area after 12 weeks of treatment. As well as a significant improvement in erythema, thickness, and scaling scores in genital lesions, guselkumab also demonstrated a favorable safety profile, with only two mild and transient treatment-related adverse events. These results suggest that the IL-23 inhibitor is an effective and safe therapeutic option for patients with genital psoriasis.<sup>49,50</sup> The management of genital psoriasis can be summarized in figure 3.

## Conclusion

Genital psoriasis is a chronic and recurrent inflammatory disease, with a possibly underestimated prevalence, which significantly impacts patients' physical, emotional, and sexual health. Its treatment involves considerable challenges; however, traditional therapies such as low-potency corticosteroids may be used as an initial approach. In addition, calcineurin inhibitors

and Vitamin D analogues are alternatives for long-term topical therapies, due to their safety profile. More recent studies involving biologic therapies, particularly IL-17A and IL-23 inhibitors, have shown promising results in the management of refractory genital psoriasis, demonstrating improvements in both clinical outcomes and quality of life.

## Funding

None.

## Conflicts of interest

None.

## Ethical considerations

**Protection of human subjects and animals.** The authors declare that no experiments on humans or animals were performed for this research.

**Confidentiality, informed consent, and ethical approval.** The authors have followed their institution's confidentiality protocols, obtained informed consent from all patients, and secured approval from the Ethics Committee. SAGER guidelines have been followed as applicable to the nature of the study.

**Declaration on the use of artificial intelligence.** The authors declare that no generative artificial intelligence was used in the writing or creation of the content of this manuscript.

## References

1. Veale DJ, Fearon U. The pathogenesis of psoriatic arthritis. *Lancet*. 2018;391:2273-84.
2. Ryan C, Sadler M, De Vol E, Patel M, Lloyd AA, Day A, et al. Genital psoriasis is associated with significant impairment in quality of life and sexual functioning. *J Am Acad Dermatol*. 2015;72:978-83.
3. Kelly A, Ryan C. Genital psoriasis: impact on quality of life and treatment options. *Am J Clin Dermatol*. 2019;20:639-46.
4. Duarte GV, Porto-Silva L, De Oliveira MF. Epidemiology and treatment of psoriasis: a Brazilian perspective. *Psoriasis (Auckl)*. 2015;5:55-64.
5. Romiti R, Amone M, Menter A, Miot HA. Prevalence of psoriasis in Brazil: a geographical survey. *Int J Dermatol*. 2017;56:e167-8.
6. Meeuwis KA, Potts Bleakman A, Van De Kerkhof PC, Dutronc Y, Hennege C, Kornberg LJ, et al. Prevalence of genital psoriasis in patients with psoriasis. *J Dermatolog Treat*. 2018;29:754-60.
7. Yi OS, Huan KY, Har LC, Ali NM, Chiang TW. Genital psoriasis: a prospective, observational, single-centre study on prevalence, clinical features, risk factors, and its impact on quality of life and sexual health. *Indian J Dermatol*. 2022;67:205.
8. Saraswat PK, Garg A, Mishra D, Garg S. Pattern of nonvenereal genital dermatoses among males attending a tertiary care center. *Indian J Sex Transm Dis AIDS*. 2014;35:129-34.
9. Mahajan R, Kumaran MS, Narang T, Handa S, Dogra S. Genital psoriasis among Indians: a prospective cross-sectional study. *Australas J Dermatol*. 2015;56:e18-20.
10. Schielein MC, Tizek L, Schuster B, Ziehfrennd S, Biedermann T, Zink A. Genital psoriasis and associated factors of sexual avoidance: a people-centered cross-sectional study in Germany. *Acta Derm Venereol*. 2020;100:adv00151.
11. Meeuwis KA, De Hullu JA, Massuger LF, Van De Kerkhof PC, Van Rossum MM. Genital psoriasis: a systematic literature review on this hidden skin disease. *Acta Derm Venereol*. 2011;91:5-11.
12. Lysell J, Tessma M, Nikamo P, Wahlgren CF, Ståhle M. Clinical characterisation at onset of childhood psoriasis - a cross sectional study in Sweden. *Acta Derm Venereol*. 2015;95:457-61.
13. Bologna JL, Schaffer JV, Cerroni L. *Dermatology*. 4th ed. Philadelphia, PA, USA: Elsevier; 2018.
14. Varghese M, Kindel S. Pigmentary disorders and inflammatory lesions of the external genitalia. *Urol Clin North Am*. 1992;19:111-21.
15. Farber EM, Nall L. Genital psoriasis. *Cutis*. 1992;50:263-6.
16. Meeuwis KA, De Hullu JA, De Jager ME, Massuger LF, Van De Kerkhof PC, Van Rossum MM. Genital psoriasis: a questionnaire-based survey on a concealed skin disease in the Netherlands. *J Eur Acad Dermatol Venereol*. 2010;24:1425-30.
17. Meeuwis KA, Van De Kerkhof PC, Massuger LF, De Hullu JA, Van Rossum MM. Patients' experience of psoriasis in the genital area. *Dermatology*. 2012;224:271-6.
18. Borghi A, Virgili A, Corazza M. Dermoscopy of inflammatory genital diseases: practical insights. *Dermatol Clin*. 2018;36:451-61.
19. Albert S, Neill S, Derrick EK, Calonje E. Psoriasis associated with vulval scarring. *Clin Exp Dermatol*. 2004;29:354-6.
20. Kapila S, Bradford J, Fischer G. Vulvar psoriasis in adults and children: a clinical audit of 194 cases and review of the literature. *J Low Genit Tract Dis*. 2012;16:364-71.
21. Czuczwar P, Stępnia A, Goren A, Wrona W, Paszkowski T, Pawlaczyk M, et al. Genital psoriasis: a hidden multidisciplinary problem - a review of literature. *Ginekol Pol*. 2016;87:717-21.
22. Butacu AI, Toma C, Negule IE, Manole I, B nic AN, Ple ea A, et al. Updates on psoriasis in special areas. *J Clin Med*. 2024;13:7549.
23. Worley B, Glassman SJ. p16+ squamous cell carcinoma *in situ* masquerading as genital psoriasis. *JAAD Case Rep*. 2017;3:542-5.
24. Jacobson CC, Kimball AB. Rethinking the psoriasis area and severity index: the impact of area should be increased. *Br J Dermatol*. 2004;151:381-7.
25. Finlay AY, Khan GK. Dermatology life quality index (DLQI). *Clin Exp Dermatol*. 1994;19:210-6.
26. Gottlieb AB, Kirby B, Ryan C, Naegeli AN, Burge R, Potts Bleakman A, et al. The development of the genital psoriasis sexual frequency questionnaire (GenPs-SFQ) to assess the impact of genital psoriasis on sexual health. *Dermatol Ther (Heidelb)*. 2018;8:33-44.
27. Gottlieb AB, Kirby B, Ryan C, Naegeli AN, Burge R, Potts Bleakman A, et al. The development of a patient-reported outcome measure for assessment of genital psoriasis symptoms: the genital psoriasis symptoms scale (GPSS). *Dermatol Ther (Heidelb)*. 2018;8:45-56.
28. Hong JJ, Mosca ML, Hader EK, Brownstone ND, Bhutani T, Liao WJ. Genital and inverse/intertriginous psoriasis: an updated review of therapies and recommendations for practical management. *Dermatol Ther (Heidelb)*. 2021;11:833-44.
29. Wozel G. Psoriasis treatment in difficult locations: scalp, nails, and intertriginous areas. *Clin Dermatol*. 2008;26:448-59.
30. Meeuwis KA, De Hullu JA, IntHout J, Hendriks IM, Sparreboom EE, Massuger LF, et al. Genital psoriasis awareness program: physical and psychological care for patients with genital psoriasis. *Acta Derm Venereol*. 2015;95:211-6.
31. Khosravi H, Siegel MP, Van Voorhees AS, Merola JF. Treatment of inverse/intertriginous psoriasis: updated guidelines from the medical board of the national psoriasis foundation. *J Drugs Dermatol*. 2017;16:760-6.
32. Reitamo S, Rissanen J, Remitz A, Granlund H, Erkkö P, Elg P, et al. Tacrolimus ointment does not affect collagen synthesis: results of a single-center randomized trial. *J Invest Dermatol*. 1998;111:396-8.
33. Bissonnette R, Nigen S, Bolduc C. Efficacy and tolerability of topical tacrolimus ointment for the treatment of male genital psoriasis. *J Cutan Med Surg*. 2008;12:230-4.
34. Rallis E, Nasiopoulou A, Kouskoukis C, Roussaki-Schulze A, Koumantaki E, Karpouzis A, et al. Successful treatment of genital and facial psoriasis with tacrolimus ointment 0.1%. *Drugs Exp Clin Res*. 2005;31:141-5.
35. Menter A, Leonardi CL, Davis DM, Gelfand JM, Lichten J, Mehta NN, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2019;80:1029-72.
36. Liao YH, Chiu HC, Tseng YS, Tsai TF. Comparison of cutaneous tolerance and efficacy of calcitriol 3 microg g(-1) ointment and tacrolimus 0.3 mg g(-1) ointment in chronic plaque psoriasis involving facial or genitofemoral areas: a double-blind, randomized controlled trial. *Br J Dermatol*. 2007;157:1005-12.
37. Stern RS. Genital tumors among men with psoriasis exposed to psoralens and ultraviolet A radiation (PUVA) and ultraviolet B radiation. The photochemotherapy follow-up study. *N Engl J Med*. 1990;322:1093-7.
38. Singh N, Thappa DM. Circinate pustular psoriasis localized to glans penis mimicking 'circinate balanitis' and responsive to dapsone. *Indian J Dermatol Venereol Leprol*. 2008;74:388-9.

39. Guglielmetti A, Conlledo R, Bedoya J, Ianiszewski F, Correa J. Inverse psoriasis involving genital skin folds: successful therapy with dapson. *Dermatol Ther (Heidelb)*. 2012;2:15.
40. Mrowietz U, Barker J, Conrad C, Jullien D, Gisondi P, Flower A, et al. Efficacy and safety of apremilast in patients with limited skin involvement, plaque psoriasis in special areas and impaired quality of life: results from the EMBRACE randomized trial. *J Eur Acad Dermatol Venereol*. 2023;37:348-55.
41. Merola JF, Parish LC, Guenther L, Lynde C, Lacour JP, Staubach P, et al. Efficacy and safety of apremilast in patients with moderate-to-severe genital psoriasis: results from DISCREET, a phase 3 randomized, double-blind, placebo-controlled trial. *J Am Acad Dermatol*. 2023;90:485-93.
42. Yosipovitch G, Foley P, Ryan C, Cather JC, Meeuwis KA, Burge R, et al. Ixekizumab improved patient-reported genital psoriasis symptoms and impact of symptoms on sexual activity vs placebo in a randomized, double-blind study. *J Sex Med*. 2018;15:1645-52.
43. Ryan C, Menter A, Guenther L, Blauvelt A, Bissonnette R, Meeuwis K, et al. Efficacy and safety of ixekizumab in a randomized, double-blinded, placebo-controlled phase IIIb study of patients with moderate-to-severe genital psoriasis. *Br J Dermatol*. 2018;179:844-52.
44. Guenther L, Potts Bleakman A, Weisman J, Poulin Y, Spelman L, Erickson J, et al. Ixekizumab results in persistent clinical improvement in moderate-to-severe genital psoriasis during a 52 week, randomized, placebo-controlled, phase 3 clinical trial. *Acta Derm Venereol*. 2020;100:adv00006.
45. Burlando M, Herzum A, Carmisciano L, Cozzani E, Parodi A. Biological therapy in genital psoriasis in women. *Dermatol Ther*. 2020;33:e13110.
46. Piaserico S, Riedl E, Pavlovsky L, Vender RB, Mert C, Tangsirirap N, et al. Comparative effectiveness of biologics for patients with moderate-to-severe psoriasis and special area involvement: week 12 results from the observational psoriasis study of health outcomes (PSoHO). *Front Med (Lausanne)*. 2023;10:1185523.
47. Carrera CG, Dapavo P, Malagoli P, Naldi L, Arancio L, Gaiani F, et al. PACE study: real-life psoriasis area and severity index (PASI) 100 response with biological agents in moderate-severe psoriasis. *J Dermatolog Treat*. 2018;29:481-6.
48. Orsini D, Gargiulo L, Ibba L, Ingurgio RC, Valenti M, Perugini C, et al. Effectiveness of risankizumab in plaque psoriasis with involvement of difficult-to-treat areas: a real-world experience from two referral centers. *J Dermatolog Treat*. 2023;34:2220849.
49. Bonifati C, Lembo S, Richetta AG, Romanelli M, Satolli F, Corazza M, et al. Effectiveness of guselkumab in patients with facial and/or genital psoriasis: interim analysis results at Week 12 from the GULLIVER study. *J Eur Acad Dermatol Venereol*. 2024;39 Suppl 1:7-14.
50. Bonifati C, Romanelli M, Corazza M, Richetta AG, Lembo S, Satolli F, et al. Effectiveness and Safety of Guselkumab: Interim Results. In: Poster Presented at: EADV Congress 2023; Oct 11-14, 2023. Berlin, Germany; 2023.